



MEETING ABSTRACT

Open Access

Effectiveness of the Tobacco Tactics program in the Department of Veterans Affairs

Sonia A Duffy^{1,2,3,4*}, David Ronis², Carrie A Karvonen-Gutierrez¹, Lee A Ewing¹, Gregory W Dalack⁴, Patricia M Smith⁵, Timothy P Carmody⁶, Thomas Hicks⁷, Christopher Hermann⁸, Pamela Reeves⁸, Petra Flanagan⁹

From 11th Annual Conference of the International Society for the Prevention of Tobacco Induced Diseases (ISPTID)

Athens, Greece. 9-11 December 2013

Background

Smoking cessation interventions during hospitalization have been shown to be efficacious, yet are rarely incorporated into practice. The purpose of this study was to determine the effectiveness of the Tobacco Tactics program in three Veterans Affairs (VA) hospitals.

Materials and methods

In this quasi-experimental pre- post- comparison effectiveness trial, inpatient nurses were educated to provide the Tobacco Tactics intervention in the Ann Arbor, MI and Detroit, MI VA hospitals, while the Indianapolis, IN VA hospital was the control site (N = 1,070). The Tobacco Tactics nurse toolkit included: 1) one contact hour for training; 2) a PowerPoint presentation on behavioral and pharmaceutical interventions; 3) a pocket card "Helping Smokers Quit: A Guide for Clinicians"; 4) pharmaceutical and behavioral protocols; and 5) a computerized template for nurse documentation. The patient toolkit included: 1) a brochure; 2) a videotape "Smoking: Getting Ready to Quit;" 3) a Tobacco Tactics manual; 4) pharmaceuticals; 5) a 1-800-QUIT-NOW help line card; and 6) post-discharge telephone calls. Smoking patients were surveyed in the hospital and again six-months post-discharge. Urinary cotinine tests were used to verify six-month smoking status.

Results

The average age was 55.3 years, most were male (94%) and not married (76%). After adjustment for the propensity of being assigned to treatment condition, there were

significant improvements in 6-month quit rates in the pre- to post-intervention time periods in Ann Arbor ($p = 0.004$) and Detroit ($p < 0.001$) compared to the Indianapolis control site. The intervention was particularly effective in Detroit where pre-intervention quit rates were 4% compared to 13% post-intervention.

Conclusions

This study showed that training staff nurses to integrate smoking cessation services into their routine care may increase quit rates. The Tobacco Tactics program, which meets the newly released (2011) Joint Commission standards that apply to all inpatient smokers, has the potential to significantly decrease smoking among patients admitted to VA hospitals.

Acknowledgements

Funding was supported by the Department of Veterans Affairs Service Directed Project (SDP 06-003).

Authors' details

¹Ann Arbor VA Center for Clinical Management Research, Health Services Research and Development, Ann Arbor, Michigan, 48105, USA. ²School of Nursing, University of Michigan, Ann Arbor, Michigan, 48105, USA. ³Department of Otolaryngology, University of Michigan, Ann Arbor, Michigan, 48105, USA. ⁴Department of Psychiatry, University of Michigan, Ann Arbor, Michigan, 48105, USA. ⁵Northern Ontario School of Medicine, Lakehead University, Sudbury, Ontario, P3E 2C6, Canada. ⁶San Francisco VA Medical Center, San Francisco, California, 94121, USA. ⁷Richard L. Roudebush VA Medical Center, Indianapolis, 46202, USA. ⁸John D. Dingell VA Medical Center, Detroit, Michigan, 48201, USA. ⁹VA Ann Arbor Healthcare System, Ann Arbor, Michigan 48105, USA.

Published: 6 June 2014

doi:10.1186/1617-9625-12-S1-A12

Cite this article as: Duffy *et al.*: Effectiveness of the Tobacco Tactics program in the Department of Veterans Affairs. *Tobacco Induced Diseases* 2014 **12**(Suppl 1):A12.

* Correspondence: bump@umich.edu

¹Ann Arbor VA Center for Clinical Management Research, Health Services Research and Development, Ann Arbor, Michigan, 48105, USA
Full list of author information is available at the end of the article