



MEETING ABSTRACT

Open Access

Design and implementation of questionnaires to assess chronic exposure to secondhand smoke in children and adults

Mary Misailidi^{1*}, Christos I Papakonstantinou¹, Manolis N Tzatzarakis², Mathaios P Kavvalakis², Yiannis Koutedakis³, Aristidis M Tsatsakis², Andreas D Flouris¹

From 11th Annual Conference of the International Society for the Prevention of Tobacco Induced Diseases (ISPTID)

Athens, Greece. 9-11 December 2013

Background

Despite a multitude of anti-smoking campaigns being active worldwide, the number of smokers is currently larger than at any other time in human history. An essential first step towards minimizing the health effects of secondhand smoke (SHS) is to accurately assess the exposure level of individuals. The objectives of this study were to: (i) to develop questionnaires that can identify never-smoking children and adults experiencing increased exposure to secondhand smoke (SHS+), (ii) to determine their validity against hair nicotine, and (iii) assess their reliability.

Materials and methods

A sample of 191 children (85 males; 106 females; 7-18 years) and 95 adult (23 males; 72 females; 18-62 years) never-smokers consented to hair nicotine analysis and answered a large number of questions pertaining to all relevant sources of SHS. A randomly-selected 30% answered the questions again after 20-30 days.

Results

Prevalence of SHS+ in children and adults was 0.52 ± 0.07 and 0.67 ± 0.10 , respectively ($p < 0.05$). The Smoke Scale for Children (SS-C) and the Smoke Scale for Adults (SS-A) were developed via factor analysis and included nine questions each. Positivity criteria for SS-C and SS-A via receiver operating characteristics (ROC) curve analysis were identified at >16.5 and >16 , respectively. Significant Kappa agreement ($p < 0.05$) was confirmed when comparing the

SS-C and SS-A to hair nicotine concentration. Reliability analyses demonstrated that the SS-C and SS-A scores obtained on two different days are highly correlated ($p < 0.001$) and not significantly different ($p > 0.05$). Area under the curve and McNemar's Chi-square showed no pair-wise differences in sensitivity and specificity at the cutoff point between the two different days for SS-C and SS-A ($p > 0.05$).

Conclusions

We conclude that the SS-C and the SS-A represent valid, reliable, practical, and inexpensive instruments to identify children and adult never-smokers exposed to increased SHS.

Authors' details

¹FAME Laboratory, Institute of Research and Technology Thessaly, Centre for Research and Technology Hellas, Trikala, 42100, Greece. ²Centre of Toxicology Science and Research, School of Medicine, University of Crete, Heraklion, 71003, Greece. ³Department of Exercise Sciences, University of Thessaly, Trikala, 42100, Greece.

Published: 6 June 2014

doi:10.1186/1617-9625-12-S1-A28

Cite this article as: Misailidi *et al.*: Design and implementation of questionnaires to assess chronic exposure to secondhand smoke in children and adults. *Tobacco Induced Diseases* 2014 **12**(Suppl 1):A28.

¹FAME Laboratory, Institute of Research and Technology Thessaly, Centre for Research and Technology Hellas, Trikala, 42100, Greece
Full list of author information is available at the end of the article